

# Chiral symmetry breaking and $\theta$ vacuum structure in QCD

G. Morchio

Dipartimento di Fisica, Università di Pisa,  
and INFN, Sezione di Pisa, Pisa, Italy

F. Strocchi

Scuola Normale Superiore, Pisa, Italy  
and INFN, Sezione di Pisa

## Abstract

The solution of the axial  $U(1)$  problem, the role of the topology of the gauge group in forcing the breaking of axial symmetry in any irreducible representation of the observable algebra and the  $\theta$  vacua structure are revisited in the temporal gauge with attention to the mathematical consistency of the derivations. Both realizations with strong and weak Gauss law are discussed; the control of the general mechanisms and structures is obtained on the basis of the localization of the (large) gauge transformations and the local generation of the chiral symmetry. The Schwinger model in the temporal gauge exactly reproduces the general results.

Keywords: Chiral symmetry; QCD; gauge group topology;  $\theta$  vacua  
PACS: 11.30.Rd; 11.30.Qc; 12.38.-t; 11.40.Ha

# 1 Introduction

The solution of the  $U(1)$  problem by the discovery of the  $\theta$  vacuum structure and its strong relation with chiral symmetry breaking has been one of the cornerstones in the theoretical analysis of the standard model of elementary particles. [2] [3] [4]

The standard (and historically the first) arguments in favor of a non-trivial role of the topology relies on semiclassical approximations, in terms of boundary conditions of the classical field configurations and their effect in the (euclidean) functional integral. These results have been excellently reviewed by Coleman, [5] [6] who also pointed out the limitation of such arguments, since smooth field configurations, in particular those of finite action, have zero functional measure. The relevance of the classification of the (smooth) field configurations in terms of their pure gauge behaviour at space infinity, and of the corresponding winding number, is therefore a non-trivial mathematical problem, especially in the infinite volume limit.

A decisive step in the direction of a mathematical control of the mechanism of  $\theta$  vacuum structure was taken by Jackiw, who emphasized the role of the topology of the gauge group, without involving the classification of the gauge field configurations and thus avoiding the problem of the semiclassical approximation of the functional integral. [7]

Jackiw's analysis is done in the temporal gauge and the very peculiar features of such a gauge raise some structural mathematical problems, with a non-trivial impact on the general argument. The aim of this note is to present an analysis which does not suffer of mathematical inconsistencies and, in a certain sense, provides a mathematical glossary to Jackiw's strategy.

The paper is organized according to the following pattern.

As in the abelian case, the invariance of the vacuum under the gauge transformations generated by the Gauss operator implies a *non-regular representation of the field algebra* (Section 2).

In Section 3, the *non-trivial topology* of the group  $\mathcal{G}$  of time independent gauge transformations is classified in terms of *group valued gauge functions  $\mathcal{U}(\mathbf{x})$  of compact support*.

This significantly simplifies the discussion with respect to the con-

ventional analysis, [7] [5] where the gauge functions are only required to have a limit, for  $|\mathbf{x}| \rightarrow \infty$ , independent of the direction (with the need of requiring for the analysis a faster than  $1/r$  decay of the gauge vector fields  $A_i^a(\mathbf{x})$ ). Furthermore, the exponentials of the topological charge (in bounded regions), at the basis of the standard analysis, are shown to have vanishing matrix elements between vectors satisfying the Gauss law constraint and therefore a non-trivial impact of the large gauge transformations on the structure of the physical states requires further ingredients.

In Section 4, it is argued that a crucial role for disclosing the physical relevance of the non-trivial topology of the gauge group is played by the fermions and the associated *chiral symmetry*.

Following Bardeen, [8] we show that, contrary to statements appeared in the literature, the presence of the chiral anomaly does not prevent the chiral symmetry from being a well defined time independent group of automorphisms of the field algebra and of its gauge invariant (observable) subalgebra, generated by the operators  $V_R^5(\lambda)$ ,  $\lambda \in \mathbf{R}$ , formally the exponentials of the charge density  $J_0(f_R \alpha_R)$ ,  $J_\mu^5$  denoting the conserved gauge dependent axial current .

The *solution of the  $U(1)$  problem* (i.e. the absence of massless Goldstone bosons associated to chiral symmetry breaking) is provided by the failure of a crucial hypotheses of the Goldstone theorem, namely the impossibility of writing the symmetry breaking Ward identities, since the vacuum expectations  $\langle J_0^5(f_R \alpha_R) \bar{\psi} \psi \rangle_0$  do not exist, as a consequence of the non-regularity of the local implementers of the chiral symmetry,  $V_R^5(\lambda)$ .

The interplay between the *topology of the gauge group* and the chiral transformations which *forces chiral symmetry breaking* (in any irreducible, or even factorial, representation of the observable algebra) and gives rise to a non-trivial vacuum structure is clearly displayed under general assumptions. By assuming that the localized large gauge transformations may be implemented by unitary operators which are “functions” of fields with the same localization, we show that the non-trivial topology of the gauge group reflects in a *non-trivial center of the algebra of observables, which is not left pointwise invariant under the chiral transformations* (Section 5).

The standard labeling of the irreducible (or even factorial) repre-

sentations of the observable algebra by an angle  $\theta \in [0, \pi)$ , ( $\theta$  sectors) is obtained by analyzing the reducible representation of the field algebra defined by a chiral invariant vacuum and by a spectral decomposition over the center of the observable algebra.

In Section 6, we consider a realization of the temporal gauge in which only a weak Gauss invariance of the vacuum is required, as discussed in the abelian case. [9] As a consequence of the lack of Gauss invariance of the vacuum, the conserved axial current  $J_\mu^5$  can be represented by a well defined field operator and Bardeen analysis directly applies.

Under the same assumption of implementation of the large gauge transformations by local operators, we show that, if the chiral symmetry is unitarily implemented, the vacuum defines a reducible representation  $\pi^0$  of the observable algebra; chiral symmetry is broken in each irreducible component. Independently of the possibility of extending the  $\theta$  vacua to non-positive weakly gauge invariant functionals on the field algebra, the chiral current  $J_\mu^5$  does not exist in the physical representation space of the observable algebra, in particular in a  $\theta$  sector.

It is worthwhile to stress that, contrary to statements appeared in the literature, the derivation of the physical consequences of the topology of  $\mathcal{G}$  crucially relies on the presence of fermions and their chiral transformations. The essential point, which distinguishes the abelian case from the non-abelian one, beyond the existence of the chiral anomaly, is the *existence of a center of the local observable algebras, following from the topology of  $\mathcal{G}$ , which is not pointwise invariant under the chiral transformations.*

All the general results are exactly reproduced by the Schwinger model in the temporal gauge, analyzed in Section 7, both in the positive non-regular realization with a Gauss invariant vacuum and in the indefinite regular (quasi free) realization, with the weak form of the Gauss law constraint.

## 2 Temporal gauge in QCD and Gauss law

For the discussion of the non-perturbative aspects of QCD, in particular the  $\theta$  vacuum structure, its relation with the topology of the gauge field configurations and its role in chiral symmetry breaking, the temporal gauge has proved to be particularly convenient.

However, as noted before for the abelian (QED) case, [9] the conflict between the Gauss law constraint and canonical quantization raises problems of mathematical consistency, which, as we shall see, affect the derivation of the general structures leading to significant physical implications.

For simplicity, we start by considering the case with only vector fields (no fermion or scalar field being present). Then, at the classical level the QCD Lagrangean density reduces to the Yang-Mills form

$$\mathcal{L} = -\frac{1}{4} \sum_a F_{\mu\nu}^a F^{\mu\nu a} = \frac{1}{2} \sum_a (\mathbf{E}_a^2 - \mathbf{B}_a^2), \quad (2.1)$$

where in the temporal gauge, defined by  $A_a^0 = 0$ ,

$$\mathbf{E}_a = -\dot{\mathbf{A}}_a, \quad \mathbf{B}_a = \nabla \times \mathbf{A}_a - \frac{1}{2} g f_{abc} \mathbf{A}_b \times \mathbf{A}_c, \quad (2.2)$$

( $a$  is a color index and  $f_{abc}$  are the structure constants of the Lie algebra of the color gauge group  $\mathcal{G}$ ).

The corresponding equations of motion, obtained by variations with respect to  $\mathbf{A}_a$ , are

$$\partial_t \mathbf{E}_a = \nabla \times \mathbf{B}_a + g f_{abc} \mathbf{A}_b \times \mathbf{B}_c \equiv (\mathbf{D} \times \mathbf{B})_a, \quad (2.3)$$

which imply

$$\partial_t G_a = 0, \quad G_a \equiv \nabla \cdot \mathbf{E}_a + g f_{abc} \mathbf{A}_b \cdot \mathbf{E}_c \equiv (\mathbf{D} \cdot \mathbf{E})_a. \quad (2.4)$$

The operators  $G_a$  are called the *Gauss law operators*.

In the standard quantum version of the temporal gauge it is assumed that the fields  $\mathbf{A}_a$  and their powers can be defined and quantization is given by the canonical commutation relations.

In particular one has the following commutation relations

$$\begin{aligned}
 -i [\mathbf{D} \cdot \mathbf{E}_a(\mathbf{x}, t), \mathbf{A}_b(\mathbf{y}, t)] &= \delta_{ab} \nabla \delta(\mathbf{x} - \mathbf{y}) + g f_{abc} \mathbf{A}_c(\mathbf{x}, t) \delta(\mathbf{x} - \mathbf{y}). \\
 -i [\mathbf{D} \cdot \mathbf{E}_a(\mathbf{x}, t), \mathbf{E}_b(\mathbf{y}, t)] &= g f_{abc} \mathbf{E}_c(\mathbf{x}, t) \delta(\mathbf{x} - \mathbf{y}).
 \end{aligned}
 \tag{2.5}$$

They state that the Gauss operators generate the infinitesimal time independent gauge transformations,  $\delta^\Lambda$ , with  $\Lambda^a(\mathbf{x}) \in \mathcal{S}(\mathbf{R}^3)$  the c-number gauge function

$$\delta^\Lambda \mathbf{A}_a(x) = \nabla \Lambda(\mathbf{x}) + g f_{abc} \mathbf{A}^b(x) \Lambda^c(\mathbf{x}). \tag{2.6}$$

Since the variables  $A_a^0$  are missing in the Lagrangean, one cannot exploit the stationarity of the action with respect to them and therefore one does not get the Gauss law  $G_a = 0$ . Actually, the Gauss law is incompatible with eq.(2.5) and therefore with canonical quantization and more crucially with the Gauss operator being the generator of the time independent gauge transformations, eq.(2.6).

A proposed solution of this conflict, widely adopted in the literature and in textbook discussions of the temporal gauge, is to require the Gauss law constraint as an operator equation on the (subspace of) physical states and, in particular, on the vacuum state. However, such a solution is not mathematically consistent. In fact, the vacuum expectation of eq.(2.6) gives zero on the left hand side and non-zero on the right hand side.

It has been proposed [7] [10] to cope with this paradox by admitting that the vacuum vector is not normalizable. In our opinion, such a solution is not acceptable, because it does not yield a representation of a field algebra containing both gauge dependent and gauge independent fields.

A mathematically acceptable solution for the Gauss law constraint is to adopt a Weyl quantization and admit non-regular representations.

As a preliminary step in this direction, we recall that from a mathematical point of view the quantum fields are operator valued distributions and a smearing with test functions (typically infinitely differentiable and of compact support) is needed for obtaining well defined Hilbert space operators. Therefore, we consider the *field algebra* generated by the polynomials of the smeared fields  $A_a^i(f)$ ,  $f \in \mathcal{S}(\mathbf{R}^4)$ ; we

shall assume that by a suitable point splitting procedure one can consider as field variables the powers of  $\mathbf{A}_a(x)$  and its derivatives, like e.g. the Gauss operator  $G_a(g)$ , the “magnetic” field  $B_a^i(g)$ ,  $g \in \mathcal{S}(\mathbf{R}^4)$ , etc.

We then take the polynomials of such field variables as a *local* (Borchers) *field algebra*  $\mathcal{F}$ , transforming covariantly under the space time translations  $\alpha_y$ ,  $y \in \mathbf{R}^4$ ,

$$\alpha_y(A_a^i(f)) = A_a^i(f_y), \quad f_y(x) = f(x - y).$$

In order to simplify the bookkeeping of the indices, it is convenient to introduce the following notations:  $T^a$  denote the hermitean representation matrices of the Lie algebra of the gauge group, normalized so that  $\text{Tr } T^a T^b = \delta_{ab}$ , the field  $A^i(x) = \sum_a A_a^i T^a$  are Lie algebra valued distributions on Lie algebra valued test functions  $\mathbf{f}^i(x) = \sum_a f_a^i(x) T^a$ ,  $f_a^i \in \mathcal{S}(\mathbf{R}^4)$ ,

$$\mathbf{A}(\mathbf{f}) \equiv \int d^4x \text{Tr} (\mathbf{A}(x) \mathbf{f}(x)) = \int d^4x \sum_{i,a} A_a^i(x) f_a^i(x).$$

Unless otherwise stated, in the following the sum over repeated space and gauge indices will be understood.

With this notation the time independent gauge transformations  $\alpha_{\mathcal{U}}$  are labeled by gauge group valued unitary  $C^\infty$  functions  $\mathcal{U}(\mathbf{x})$ , which may be taken to differ from the identity only on a compact set  $K_{\mathcal{U}}$ , and

$$\alpha_{\mathcal{U}}(\mathbf{A}(\mathbf{f})) = \mathbf{A}(\mathcal{U}\mathbf{f}\mathcal{U}^{-1}) + \mathcal{U}\partial\mathcal{U}^{-1}(\mathbf{f}), \quad (2.7)$$

$$\mathcal{U}\partial\mathcal{U}^{-1}(\mathbf{f}) \equiv \int d^4x \text{Tr} \left( \sum_i \mathcal{U}(\mathbf{x}) \partial^i \mathcal{U}^{-1}(\mathbf{x}) \mathbf{f}^i(x) \right).$$

The (space-time) localization of  $\mathcal{U}$  is given by the cylinder  $C_{\mathcal{U}} \equiv K_{\mathcal{U}} \times \mathbf{R}$ , so that  $\alpha_{\mathcal{U}}(\mathbf{A}(\mathbf{f})) = \mathbf{A}(\mathbf{f})$  if  $\text{supp } \mathbf{f} \cap C_{\mathcal{U}} = \emptyset$ .

We denote by  $\mathcal{U}^\lambda$ ,  $\lambda \in \mathbf{R}$ , the gauge functions corresponding to one-parameter subgroups of the gauge group. They can be written in the form  $\mathcal{U}^\lambda(\mathbf{x}) = e^{i\lambda g(\mathbf{x})}$  with  $g = \sum_a g_a(\mathbf{x}) T^a$  a Lie algebra valued function, infinitely differentiable and of compact support ( $g_a \in \mathcal{D}(\mathbf{R}^3)$ ). All gauge transformations of compact support in a neighborhood of the identity, in the  $C^\infty$  topology, are of this form; they generate the Gauss subgroup  $\mathcal{G}_0$  of the gauge group  $\mathcal{G}$ .

For discussing the Weyl quantization one has to consider the *exponential field algebra*  $\mathcal{F}_W$  generated by the unitary operators  $W(\mathbf{f})$ ,  $\mathbf{f}^i = \sum_a f_a^i(x) T^a$ ,  $f_a^i \in \mathcal{S}(\mathbf{R}^4)$ , formally the exponentials  $e^{i\mathbf{A}(\mathbf{f})}$ , and by unitary operators  $V(\mathcal{U}^\lambda)$ , representing  $\mathcal{G}_0$ , formally the exponentials of the Gauss operators,

$$V(\mathcal{U}^\lambda) = e^{i\lambda G(g)}, \quad G(g) = \sum_a G_a(g_a), \quad g_a \in \mathcal{D}(\mathbf{R}^3),$$

transforming covariantly under space translations. Their time independence formally follows if the dynamics  $\alpha_t$  is generated by local gauge invariant Hamiltonians  $H_R$ , so that

$$(d/dt)\alpha_t(V(\mathcal{U}^\lambda)) = i \lim_{R \rightarrow \infty} [H_R, V(\mathcal{U}^\lambda)] = 0. \quad (2.8)$$

The *observable field subalgebra* of  $\mathcal{F}_W$  is characterized by its point-wise invariance under gauge transformations.

A representation of  $\mathcal{F}_W$  also defines a representation of  $\mathcal{F}$  only if it is **regular**, i.e. if (the representatives of) the field exponentials  $W(\lambda \mathbf{f})$ ,  $\lambda \in \mathbf{R}$  define weakly continuous one-parameter groups.

A state  $\omega$  on the exponential field algebra  $\mathcal{F}_W$ , in particular a vacuum state, is said to satisfy the **Gauss law** in exponential form, if  $\omega(V(\mathcal{U}^\lambda)) = 1$ , equivalently if its representative vector  $\Psi_\omega$  in the GNS Hilbert space  $\mathcal{H}_\omega$  (defined by the expectations of  $\mathcal{F}_W$  on  $\omega$ ) satisfies

$$V(\mathcal{U}^\lambda) \Psi_\omega = \Psi_\omega, \quad \forall \mathcal{U}^\lambda. \quad (2.9)$$

Briefly, a vector state  $\Psi$  satisfying eq. (2.9) is said to be **Gauss invariant**. An operator in  $\mathcal{H}$  is *Gauss invariant* if it commutes with all the  $V(\mathcal{U}^\lambda)$ .

In the following we shall consider the realization of the temporal gauge defined by a vacuum state  $\omega$  satisfying the Gauss law. As in other interesting quantum mechanical models, like the electron in a periodic potential (Bloch electron), the quantum particle on a circle, the Quantum Hall electron, etc., the invariance of the ground state under a group of gauge transformations implies that the corresponding representation of the exponential field algebra is not regular. [11]



**Proposition 2.1** *A vacuum state  $\omega$  on the exponential field algebra  $\mathcal{F}_W$ , satisfying the Gauss law, defines a non-regular representation of  $\mathcal{F}_W$ , since :*

$$\begin{aligned}\omega(W(\mathbf{f}^i)) &= 0, & \text{if } \mathbf{f}^i(x) \neq 0, \\ &= 1, & \text{if } \mathbf{f} = 0.\end{aligned}\tag{2.10}$$

*The fields  $\mathbf{A}$ , formally the generators of the  $W(\mathbf{f})$ , cannot be defined in the GNS Hilbert space defined by the vacuum expectations and in particular the two point function of the gauge potential does not exist, only (the vacuum expectations of) the exponential functions (and of course the gauge invariant functions) of  $\mathbf{A}$  can be defined.*

*In the free case, i.e. for vanishing gauge coupling constant, the exponential field algebra becomes a Weyl field algebra, generated by the exponentials of  $\mathbf{A}_a$  and of its conjugate momenta  $\mathbf{E}_a$ , and eqs. (2.10) uniquely determine its representation as a **non-regular Weyl quantization**.*

**Proof.** For each  $\mathbf{f}^i$  there is a one-parameter subgroup  $\mathcal{U}^\lambda$  such that  $\mathcal{U}^\lambda \mathbf{f}^i \mathcal{U}^{-\lambda} = \mathbf{f}^i$ , and  $\exp iD_{\mathcal{U}^\lambda}(\mathbf{f}^i) \equiv \exp(i\mathcal{U}^\lambda \partial \mathcal{U}^{-\lambda}(\mathbf{f}^i)) \neq 1$ ; therefore, by eq. (2.9), one has

$$\omega(W(\mathbf{f}^i)) = \omega(V(\mathcal{U}^\lambda) W(\mathbf{f}^i) V(\mathcal{U}^\lambda)^*) = e^{iD_{\mathcal{U}^\lambda}(\mathbf{f}^i)} \omega(W(\mathbf{f}^i)),$$

and eqs. (2.10) follow.

Clearly, the one-parameter groups defined by  $W(\mathbf{f})$  cannot be weakly continuous and therefore the corresponding generators, i.e. the fields  $A_a^i(f)$  do not exist as operators in the GNS Hilbert space defined by the expectations of  $\mathcal{F}_W$  on  $\omega$ . The free case can be worked out along the same lines as for the abelian case. [9]

The Hilbert space  $\mathcal{H}$  of the representation of  $\mathcal{F}_W$  defined by the vacuum state  $\omega$ , satisfying the Gauss law, contains a subspace  $\mathcal{H}'$  of Gauss invariant vectors

$$V(\mathcal{U}^\lambda) \Psi = \Psi, \quad \forall \Psi \in \mathcal{H}', \quad \forall \mathcal{U}^\lambda.$$

We denote by  $\pi_0$  the representation of  $\mathcal{G}_0$  in  $\mathcal{H}$  given by the  $V(\mathcal{U}^\lambda)$ .

It is worthwhile to remark that the local operator  $v(\mathcal{U}^\lambda) \equiv V(\mathcal{U}^\lambda) - 1$  is non zero in  $\mathcal{H}$ , since  $V(\mathcal{U}^\lambda)$  implements the time independent gauge

transformations, corresponding to the one-parameter subgroups, which are non-trivial on  $\mathcal{F}_W$ . Thus, the assumptions of the Reeh-Schlieder theorem, according to which a local operator which annihilates the vacuum must vanish, cannot be satisfied.

One can easily check that the crucial point in the proof of the theorem fails, namely  $\mathcal{F}_W(\mathcal{O}) \Psi_0$ , where  $\mathcal{F}_W(\mathcal{O})$  is the exponential field algebra localized in the region  $\mathcal{O}$ , is not dense in  $\mathcal{H}$ . In fact, for any  $\mathcal{O}$  disjoint from  $C_{\mathcal{U}^\lambda}$ , by locality one has

$$(v(\mathcal{U}^\lambda) \mathcal{F}_W \Psi_0, \mathcal{F}_W(\mathcal{O}) \Psi_0) = (\mathcal{F}_W \Psi_0, \mathcal{F}_W(\mathcal{O}) v(\mathcal{U}^\lambda)^* \Psi_0) = 0,$$

$$v(\mathcal{U}^\lambda) \mathcal{F}_W \Psi_0 \neq 0.$$

Actually, the Reeh-Schlieder theorem does not apply because the relativistic spectral condition fails in  $\mathcal{H}$ . In fact, the implementers  $U(\mathbf{a})$  of the space translations are not weakly continuous, since, as in the proof of Proposition 2.1,

$$\begin{aligned} \omega(W(\mathbf{f})U(\mathbf{a})W(-\mathbf{f})) &= \omega(W(\mathbf{f})W(-\mathbf{f}_\mathbf{a})) = \\ &= e^{iD_{\mathcal{U}^\lambda}(\mathbf{f}-\mathbf{f}_\mathbf{a})} \omega(W(\mathbf{f})W(-\mathbf{f}_\mathbf{a})), \end{aligned}$$

so that the right hand side vanishes if  $\mathbf{a} \neq 0$  and it is  $= 1$ , otherwise. The spectral condition for the Fourier transforms of the matrix elements of  $U(\mathbf{a}, t) = U(\mathbf{a})U(t)$  is violated if there is strong continuity in  $t$ , since it would imply that, after smearing in time, the Fourier transform in  $\mathbf{a}$  is finite measure and therefore continuity in  $\mathbf{a}$  of the above matrix elements.

It is worthwhile to remark that the one-parameter groups  $V(\mathcal{U}^\lambda)$  are not assumed to be weakly continuous in  $\lambda$ ; actually, continuity cannot hold if the global gauge group is simple and has rank at least two (as in the case of color  $SU(3)$ ), since then one obtains the vanishing of  $\omega(W(\mathbf{f})V(U_\lambda)W(-\mathbf{f}))$ , for  $\lambda \neq 0$ ,  $f_c^i(x) = \delta_{ca}f^i(x)$ ,  $\mathcal{U}^\lambda \mathbf{f}^i \mathcal{U}^{-\lambda} = f^i(x)T^b$ ,  $[T^a, T^b] = 0$ , from the invariance of  $\omega$  under the subgroup generated by  $T^a$ , as in the proof of Proposition 2.1. Thus, in this case the Gauss law constraint can only be imposed in the exponential form.

### 3 Topology of the gauge group

By definition, eq. (2.7), we consider localized gauge functions; they obviously extend to the one point compactification of  $\mathbf{R}^3$ ,  $\dot{\mathbf{R}}^3$ , which is isomorphic to the three-sphere  $S^3$ ,

$$\mathcal{U}(\mathbf{x}) : \dot{\mathbf{R}}^3 \sim S^3 \rightarrow \mathcal{G}.$$

Such maps fall into disjoint homotopy classes labeled by “winding” numbers  $n$

$$n(\mathcal{U}) = (24\pi^2)^{-1} \int d^3x \varepsilon^{ijk} \text{Tr} (\mathcal{U}_i(\mathbf{x}) \mathcal{U}_j(\mathbf{x}) \mathcal{U}_k(\mathbf{x})) \equiv \int d^3x n_{\mathcal{U}}(\mathbf{x}),$$

where  $\mathcal{U}_i(\mathbf{x}) \equiv \mathcal{U}(\mathbf{x})^{-1} \partial_i \mathcal{U}(\mathbf{x})$ .

The gauge transformations with  $n \neq 0$  are called *large gauge transformations*. Those with zero winding number are called *small*; since they are contractible to the identity, they are products of  $\mathcal{U}(\mathbf{x})$  which are close to the identity (in the  $C^\infty$  topology) and therefore are expressible as products of  $\mathcal{U}^\lambda$ , i.e. they are elements of  $\mathcal{G}_0$ . Clearly, all elements of  $\mathcal{G}_0$  have zero winding number.

The following analysis may be also applied to gauge transformations  $\mathcal{U}(\mathbf{x})$  which are only required to have a limit for  $|\mathbf{x}| \rightarrow \infty$ , since they are of compact support modulo Gauss transformations  $\mathcal{U}^\lambda$  with such a behaviour; only the existence of the corresponding Gauss operators  $V(\mathcal{U}^\lambda)$  is required, with no additional implications.

In the above realization of the temporal gauge, the small gauge transformations are implemented by the unitary operators  $V(\mathcal{U}^\lambda) \in \mathcal{F}_W$ ; the next question is the implementability of the large gauge transformations and their distinction from the small on the Gauss invariant states. In fact, the non-triviality of the large gauge transformations on the physical space turns out to be a rather subtle question as displayed by the following Proposition.

**Proposition 3.1** *A local field operator invariant under Gauss gauge transformations is also invariant under large gauge transformations.*

*Any vector  $\Psi \in \mathcal{H}'$ , in particular the vacuum vector, defines a state, i.e. expectations, on  $\mathcal{F}_W$  invariant also under the large gauge transformations.*

**Proof.** In fact, given a gauge function  $\mathcal{U}_n(\mathbf{x})$  and its space translated by  $\mathbf{a}$ ,  $\mathcal{U}_n^a(\mathbf{x}) = \mathcal{U}_n(\mathbf{x} - \mathbf{a})$ , the combined gauge transformations  $\alpha_{\mathcal{U}_n^a}^{-1} \alpha_{\mathcal{U}_n}$  and  $\alpha_{\mathcal{U}_n} \alpha_{\mathcal{U}_n^a}^{-1}$  have zero winding number and therefore are small gauge transformations, say  $\alpha_{\mathcal{U}_0}$  and  $\alpha_{\mathcal{U}'_0}$  respectively. Then, for any local field operator  $F$  invariant under small gauge transformations one has

$$\alpha_{\mathcal{U}_n}(F) = \alpha_{\mathcal{U}_n^a} \alpha_{\mathcal{U}_0}(F) = \alpha_{\mathcal{U}_n^a}(F),$$

and for  $|\mathbf{a}|$  sufficiently large, by locality  $\alpha_{\mathcal{U}_n^a}(F) = F$ .

Quite generally, for any (local) operator  $F \in \mathcal{F}_W$  one has, for  $|\mathbf{a}|$  sufficiently large,

$$\alpha_{\mathcal{U}_n}(F) = \alpha_{\mathcal{U}'_0} \alpha_{\mathcal{U}_n^a}(F) = \alpha_{\mathcal{U}'_0}(F);$$

by the Gauss invariance of  $\omega$ , this implies  $\omega(\alpha_{\mathcal{U}_n}(F)) = \omega(F)$ .

By a standard argument, the invariance of the vacuum under the large gauge transformations implies that the gauge transformations are implemented by unitary operators; they can be chosen to represent the gauge group  $\mathcal{G}$ , to coincide the  $V(\mathcal{U}^\lambda)$ , for all Gauss transformations, and to transform covariantly under space translations.

Furthermore, if, as we assume, the dynamics is generated by local gauge invariant Hamiltonians, the implementers commute with the time translations. The implementers are unique, up to phases, if the field algebra is irreducible in  $\mathcal{H}$ ; in this case the implementers of the large gauge transformations are multiples of the identity in  $\mathcal{H}'$ .

The above results indicate that the distinction between the small and the large transformations at the level of physical states is problematic and a crucial question is the implication, if any, of the non-trivial topology of the gauge group on the physical states, more generally on the representations of the observable algebra.

One of the standard (and historically the first) arguments in favor of a non-trivial role of the topology of the gauge transformations relies on semiclassical approximations, in terms of boundary conditions of the classical field configurations and their effect on the (euclidean) functional integral. This approach has been excellently reviewed by Coleman, [5] who also pointed out the limitation of such arguments, since smooth field configurations, in particular those of finite action, have zero functional measure.

The relevance of the classification of the (smooth) field configurations in terms of their pure gauge behaviour at space infinity, and of the corresponding winding number, is therefore a non-trivial mathematical problem, especially in the infinite volume limit. The topological classification is done in finite volume, but in the infinite volume limit a non-trivial instanton density implies, in the dilute gas approximation, that the functional measure is concentrated on configurations with divergent topological number. The association of the non-trivial topology with the existence of instanton solutions, which minimize the classical action and as such do not have compact support, is probably the reason why localized large gauge transformations have not been considered in the literature.

As we shall see below, the presence of fermions plays a crucial role for the non-trivial effects of the topology of the gauge group.

Another argument for the physical consequences of the gauge group topology has been proposed in terms of the topological charge. [7] The so-called *topological current* is formally defined by

$$C^\mu(x) = -(16\pi^2)^{-1} \varepsilon^{\mu\nu\rho\sigma} \text{Tr}(F_{\nu\rho}(x) A_\sigma(x) - \frac{2}{3} A_\nu(x) A_\rho(x) A_\sigma(x)). \quad (3.1)$$

$$\partial_\mu C^\mu(x) = -(16\pi^2)^{-1} \text{Tr}^* F_{\mu\nu}(x) F_{\mu\nu}(x) \equiv \mathcal{P},$$

where  $A_\mu = (0, A_i)$ ,  $^*F_{\mu\nu} \equiv \varepsilon_{\mu\nu\rho\sigma} F^{\rho\sigma}$ . In the mathematical literature, for classical fields,  $\mathcal{P}$  is called the ‘‘Pontryagin density’’ and  $C_\mu$  the ‘‘Chern-Simons secondary characteristic class’’.

At the classical level, one has [7] the following transformation law of  $C_0(x)$  under gauge transformations  $\alpha_{\mathcal{U}}$ , defined by eqs. (2.7),

$$\alpha_{\mathcal{U}}(C_0(x)) = C_0(x) - (8\pi^2)^{-1} \partial_i [\varepsilon^{ijk} \text{Tr}(\partial_j \mathcal{U}(\mathbf{x}) \mathcal{U}(\mathbf{x})^{-1} A_k(x))] + n_{\mathcal{U}}(\mathbf{x}). \quad (3.2)$$

Therefore, at the classical level the space integral of  $C_0(\mathbf{x}, x)$  is invariant under small transformations, but it gets shifted by  $n$  under gauge transformations with winding number  $n$ .

For the quantum case one meets non-trivial consistency problems. First of all the formal expression in the right hand side of eq. (3.3) requires a point splitting regularization. It is reasonable to assume that this can be done by keeping the transformation properties of the formal expression under large gauge transformations, eq. (3.4).

The next problem is the space integral of  $C_0(x)$ . The space integrals of charge densities, even for conserved currents, are known to diverge and suitable regularizations are needed, including a time smearing (see eq. (3.5) below). In the case of conserved currents, under some general conditions one may obtain the convergence of a suitably regularized integral of the charge density, in matrix elements on states with suitable localization; [12] [13] [14] [15] but in the general case the problem seems to be open.

Actually, in the quantum temporal gauge an even more serious problem arises by the gauge dependence of  $C_\mu(x)$ . By Proposition 2.1, the regularized space integral of  $C_0(x)$

$$C_0(f_R\alpha_R) \equiv \int d^4x f_R(\mathbf{x}) \alpha_R(x_0) C_0(x), \quad (3.3)$$

where  $f_R(\mathbf{x}) = f(|\mathbf{x}|/R)$ ,  $f(x) = 1$ , for  $|x| \leq 1$ ,  $= 0$ , for  $|x| \geq 1 + \varepsilon$ ,  $\alpha_R(x_0) = \alpha(x_0/R)/R$ ,  $\int dx_0 \alpha(x_0) = \tilde{\alpha}(0) = 1$ , cannot exist as an operator in  $\mathcal{H}$ , only its exponential  $V^C(f_R\alpha_R)$ , formally  $\exp[i C_0(f_R\alpha_R)]$ , may be defined. Furthermore, as shown by the following Proposition, such exponentials have vanishing expectation on Gauss invariant states, i.e. their restriction to the physical states vanishes.

**Proposition 3.2** *The operators  $V^C(\lambda f_R\alpha_R)$ , formally the exponentials  $\exp[i\lambda C_0(f_R\alpha_R)]$  of the regularized space integrals  $C_0(f_R\alpha_R)$  of  $C_0(x)$ , and therefore assumed to transform under gauge transformations as such exponentials, cannot be weakly continuous in  $\lambda$  and therefore the field  $C_0(f_R\alpha_R)$  cannot be defined.*

*Furthermore, the  $V^C(f_R\alpha_R)$  satisfy, for all Gauss invariant vectors  $\Psi, \Phi$ ,*

$$(\Psi, V^C(f_R\alpha_R) \Phi) = 0. \quad (3.4)$$

**Proof.** In fact, if  $C_0(f)$ ,  $f \in \mathcal{D}(\mathbf{R}^4)$ , exists, by using the Gauss gauge invariance of the vacuum state  $\omega$ , the vanishing of  $\omega(A_k)$  by rotational invariance and eq. (3.4), one has

$$\begin{aligned} \omega(C_0(f)) &= \omega(V(\mathcal{U}^\lambda) C_0(f) V(\mathcal{U}^\lambda)^{-1}) = \\ &= \omega(C_0(f)) + \int d^3x f(\mathbf{x}, t) n_{\mathcal{U}^\lambda}(\mathbf{x}). \end{aligned}$$

Since for any  $f$  there is at least one  $\mathcal{U}^\lambda(\mathbf{x})$  such that the last term on the right hand side does not vanish, one gets a contradiction. Thus, only the exponential of  $C_0(f)$  can be defined.

Moreover, given  $f_R$  one can find a small gauge transformation  $\mathcal{U}(\mathbf{x})$ , with

$$\begin{aligned}\mathcal{U}(\mathbf{x}) &= \mathcal{U}_1(\mathbf{x})\mathcal{U}_2(\mathbf{x}), \quad n_{\mathcal{U}_1} + n_{\mathcal{U}_2} = 0, \quad n_{\mathcal{U}_1} \neq 0, \\ f_R \mathcal{U}_2 &= \mathbf{1}, \quad f_R \mathcal{U}_1 = \mathcal{U}_1.\end{aligned}$$

Then,  $\partial_i f_R \partial_j \mathcal{U} = 0$  and the second term on the right hand side of eq. (3.4) vanishes; furthermore  $\int d^3x n_{\mathcal{U}}(\mathbf{x}) f_R(\mathbf{x}) = n_{\mathcal{U}_1}$ . Hence, one has

$$\begin{aligned}(\Psi, V^C(f_R \alpha_R) \Phi) &= (\Psi, V(\mathcal{U}) V^C(f_R \alpha_R) V(\mathcal{U})^{-1} \Phi) = \\ &= e^{in_{\mathcal{U}_1}} (\Psi, V^C(f_R \alpha_R) \Phi)\end{aligned}$$

and eq. (3.4) follows.

In conclusion, one cannot directly exploit the non-invariance of  $C_0(x)$  for proving a non-trivial action of the large gauge transformations in  $\mathcal{H}'$ ; it is essential to take into account the non regularity of  $\exp i C_0(f_R \alpha_R)$ , its non observability and the non-existence of the limit  $R \rightarrow \infty$ .

## 4 Chiral symmetry and solution of the $U(1)$ problem

The situation changes substantially in the presence of massless fermions, since the role of the topological current is taken by a *conserved* current; hence, there is a symmetry associated to it and the crucial point is its relations with the implementers of the large gauge transformations.

In this case, the Lagrangean, eq. (2.1) gets modified by the addition of the (gauge invariant) fermion Lagrangean and the Gauss operators become

$$G_a = (\mathbf{D} \cdot \mathbf{E})_a - j_0^a, \quad j_\mu^a = ig \bar{\psi} \gamma_\mu t^a \psi.$$

The time independent gauge transformations of the fermion fields in the fundamental representation of the gauge group are

$$\alpha_{\mathcal{U}}(\psi(x)) = \mathcal{U}(\mathbf{x})\psi(x).$$

At the classical level, the Lagrangean is invariant under the one-parameter group of *chiral transformations*  $\beta^\lambda$ ,  $\lambda \in \mathbf{R}$ ,

$$\beta^\lambda(\psi) = e^{\lambda\gamma_5} \psi, \quad \beta^\lambda(\bar{\psi}) = \bar{\psi} e^{\lambda\gamma_5}, \quad \gamma_5^* = -\gamma_5, \quad \beta^\lambda(\mathbf{A}) = \mathbf{A}.$$

Correspondingly, there is a conserved current  $j_\mu^5 = ig\bar{\psi}\gamma^5\gamma_\mu\psi$ , the gauge invariant fermion axial current.

In the quantum case, a gauge invariant point splitting regularization is needed for the definition of  $j_\mu^5$  and this inevitably leads to an *anomaly*,

$$\partial^\mu j_\mu^5 = -2\partial^\mu C_\mu = -2\mathcal{P}.$$

The conserved axial current is now the gauge dependent current

$$J_\mu^5(x) = j_\mu^5(x) + 2C_\mu,$$

its conservation being equivalent to the anomaly equation for  $j_\mu^5$ .

For the discussion of the Weyl quantization, we take as local exponential field algebra  $\mathcal{F}_W$  the algebra generated by the operators  $W(\mathbf{f})$ , by the gauge invariant bilinear functions of the fermion fields and by the unitary operators  $V^5(f)$ ,  $f \in \mathcal{S}(\mathbf{R}^4)$ , formally the exponential of  $J_0^5(f)$ .

As shown by Bardeen [8] on the basis of perturbative renormalization in local gauges, the above (time independent) chiral transformations of the fermion fields are generated by the quantum field operator  $J_\mu^5(x)$  and not by the gauge invariant non conserved current  $j_\mu^5$ ; the continuity equation of  $J_\mu^5$  plays a crucial role in Bardeen analysis.

This justifies our assumption that the exponential field algebra contains the (formally defined) exponential of the smeared field  $J_\mu^5$  and in particular of the regularized integral of the charge density  $J_0^5(f_R\alpha_R)$ ; the corresponding one-parameter group of unitary operators is denoted by  $V_R^5(\lambda) = V^5(\lambda f_R\alpha_R)$ .

With the same motivations given before, eqs. (2.7) are assumed to hold together with the following transformation law of  $V_R^5(\lambda)$  under gauge transformations: for  $R$  large enough so that  $f_R(\mathbf{x}) = 1$  on the localization region of  $\mathcal{U}_n(\mathbf{x})$ , one has

$$\alpha_{\mathcal{U}_n}(V_R^5(\lambda)) = e^{i2n\lambda} V_R^5(\lambda). \quad (4.1)$$



This relation formally reflects the transformation properties of  $C_0(f)$  and the gauge invariance of  $j_0^5$ . By the proof of Proposition 3.3,  $V_R^5(\lambda)$  is not weakly continuous in  $\lambda$  and its formal generator  $J_0^5(f_R\alpha_R)$  does not exist. However,  $V_R^5(\lambda)$  act as local implementers of  $\beta^\lambda$ : in fact, since  $J_\mu^5$  is conserved, most of the standard wisdom is available [16] and one has

$$\lim_{R \rightarrow \infty} V_R^5(\lambda) F V_R^5(-\lambda) = \beta^\lambda(F), \quad \forall F \in \mathcal{F}_W. \quad (4.2)$$

It is important to stress that, thanks to locality, the above limit is reached for finite values of  $R$ , and that it preserves locality and gauge invariance. Thus, contrary to what is stated in the literature, the presence of the chiral anomaly does not prevent the chiral symmetry from being a well defined time independent automorphism of the field algebra  $\mathcal{F}_W$  of its gauge invariant (observable) subalgebra.

The loss of chiral symmetry is therefore a genuine phenomenon of spontaneous symmetry breaking and the confrontation with the Goldstone theorem becomes a crucial issue, the so-called **U(1) problem**.

The absence of parity doublets requires that the chiral symmetry be broken and the  $U(1)$  problem amounts to explaining the absence of the corresponding Goldstone massless bosons.

As discussed above, one of the basic assumptions of the Goldstone theorem, namely the existence of an automorphism of the algebra of observables, which commutes with space and time translations is satisfied.

The second crucial property, needed the proof of the theorem, is the local generation of the symmetry by a conserved current at least in expectations on the vacuum state, i.e.

$$\frac{d}{d\lambda} \langle \beta^\lambda(A) \rangle_{\lambda=0} = \langle \delta^5(A) \rangle = i \lim_{R \rightarrow \infty} \langle [J_0^5(f_R\alpha_R), A] \rangle. \quad (4.3)$$

Since the chiral automorphism  $\beta^\lambda$  is  $C^\infty$  in  $\lambda$  its generator  $\delta^5$  is well defined, but the problem is its relation with the formal generator of the unitary one-parameter group defined by the  $V_R^5(\lambda)$ .

As a matter of fact, even if  $\beta^\lambda$  can be described by the action of the local operators  $V_R^5(\lambda)$ , eq. (4.2), the non-regularity of the one-parameter unitary group  $V_R^5(\lambda)$ , prevents the existence of the corresponding generator  $J_0^5(f_R\alpha_R)$ , so that one cannot write the symmetry breaking Ward identities and obtain the Goldstone energy-momentum spectrum.

Quite generally, one has

**Proposition 4.1** *If  $\omega$  is a gauge invariant vacuum state and  $A$  is an observable symmetry breaking order parameter, i.e.*

$$\omega(\beta^\lambda(A)) \neq \omega(A) \neq 0,$$

*(the standard candidate being  $\bar{\psi}\psi$ ) then the expectations  $\omega(J_0^5(f_R\alpha_R) A)$  cannot be defined and eq. (4.3) does not hold.*

**Proof.** In fact, otherwise, for  $R$  sufficiently large, one would have

$$\omega(J_0^5(f_R\alpha_R) A) = \omega(\alpha_{\mathcal{U}_n}(J_0^5(f_R\alpha_R) A)) = \omega(J_0^5(f_R\alpha_R) A) + 2n \omega(A), \quad (4.4)$$

i.e. a contradiction.

Clearly, by Proposition 3.1, the above Proposition applies to the Gauss invariant state  $\Psi_0$ . The impossibility of writing expectations involving  $J_\mu^5$  on a gauge invariant vacuum state, solves the problems raised by R.J. Crewther in his analysis of chiral Ward identities. [17]

It is worthwhile to remark that, for the evasion of the Goldstone theorem discussed above, the occurrence of the so-called chiral anomaly (which is present also in the abelian case) is not enough; the crucial ingredient is eq. (4.1), which directly implies the non-regularity of the unitary operators  $V_R^5(\lambda)$  and the non-existence of the local charges  $J_0^5(f_R\alpha_R)$  in expectations on a gauge invariant vacuum state.

## 5 Topology, chiral symmetry breaking and vacuum structure

The implications of the non-trivial topology of the gauge group on the breaking of the chiral symmetry crucially involve the chiral transformations of the implementers of gauge group. Such chiral transformations do not follow merely from eq. (4.1), because eq. (4.2) applies to the elements of the local field algebra and its extension to implementers, even if they are strong limits of elements of  $\mathcal{F}_W$ , is not uniquely defined. This problem does not arise if there are implementers  $V(\mathcal{U}_n)$  belonging to the local field algebra.

Under this assumption, we shall first prove that the chiral symmetry is broken in any irreducible (or factorial) representation of the field algebra, as well as in any factorial representation of the observable algebra in the Gauss invariant subspace  $\mathcal{H}'$  (we recall that a representation is factorial if the center of the strong closure consists of multiples of the identity there).

We shall later show that the standard vacuum structure is obtained by analyzing the decomposition of  $\mathcal{H}'$  into factorial representation of  $\mathcal{A}$  ( $\theta$  sectors), in the case of a reducible representation of the field algebra defined by a chiral invariant vacuum; a crucial ingredient for the derivation are the non-trivial chiral transformations of the implementers  $V(\mathcal{U}_n)$ , which uniquely follow from eqs. (4.1), (4.2), since they belong to the field algebra.

The local structure of the gauge transformations imply that the implementers  $V(\mathcal{U})$  commute with the fields localized in spacetime regions  $\mathcal{O}$  disjoint from the spacetime localization region  $C_{\mathcal{U}}$  of  $\mathcal{U}$ ; thus the  $V(\mathcal{U})$  are local with respect to the field algebra, with localization region of  $V(\mathcal{U})$  given by  $C_{\mathcal{U}}$ . This locality property partly motivate the assumption of existence of *local* implementers, in the following sense.

As it is standard, in the following the local field algebras  $\mathcal{F}_W(\mathcal{O})$  and their gauge invariant (“observable”) subalgebras  $\mathcal{A}(\mathcal{O})$  are taken as strongly closed;  $\mathcal{F}_W$  and  $\mathcal{A}$  will denote their unions over  $\mathcal{O}$ . For any bounded region  $\mathcal{O}$ , the local algebra  $\mathcal{F}_W(\mathcal{O})$  may be identified with the strong closure of the polynomial algebra generated by the field exponentials  $W(\mathbf{f})$ , by the gauge invariant bilinear functions of the fermion fields and by the unitary operators  $V^5(f)$ ; as it is standard, the center of  $\mathcal{F}_W(\mathcal{O})$  is assumed to be trivial.

Then, we assume that if  $\text{supp } \mathcal{U}$  is localized in  $\mathcal{O}$ , the gauge transformation  $\alpha_{\mathcal{U}}$  may be implemented by a unitary operator  $V(\mathcal{U}) \in \mathcal{F}_W(\mathcal{O})$ . [18] Since the center of  $\mathcal{F}_W(\mathcal{O})$  is trivial, such a local implementer is unique; in particular, for Gauss transformations  $\mathcal{U}^\lambda$  it reduces to the previously introduced Gauss operators.

Then, we have

**Proposition 5.1** *The local implementers  $V(\mathcal{U})$  of the gauge transformations are of the form  $V(\mathcal{U}) F \Psi_0 = \alpha_{\mathcal{U}}(F) C_{\mathcal{U}} \Psi_0$ , where  $C_{\mathcal{U}}$  commutes with  $\mathcal{F}_W$  and belongs to the strong closure of  $\mathcal{F}_W$ .  $C_{\mathcal{U}_n}$  depends*

only on  $n$  and on the Gauss invariant vectors one has

$$V(\mathcal{U}_n) \Psi = C_n \Psi, \quad \forall \Psi \in \mathcal{H}'. \quad (5.1)$$

The algebra generated by the  $C_n$  is abelian and one has

$$C_n C_m = C_{n+m}. \quad (5.2)$$

Furthermore, one has

i) for any  $\mathcal{O}$ , there exist local operators  $C_n(\mathcal{O})$  belonging to the local observable algebra  $\mathcal{A}(\mathcal{O})$ , satisfying  $[C_n(\mathcal{O}), \mathcal{A}] = 0$  and

$$C_n(\mathcal{O}) C_m(\mathcal{O}) = C_{n+m}(\mathcal{O}), \quad C_n(\mathcal{O}) \Psi = C_n \Psi, \quad \forall \Psi \in \mathcal{H}', \quad \forall \mathcal{O}, \quad (5.3)$$

ii)  $s - \lim_{\mathcal{O} \rightarrow \mathbf{R}^4} C_n(\mathcal{O}) = C_n P_0$ ,  $P_0 \mathcal{H} = \mathcal{H}'$ .

**Proof.** In fact, if  $F \in \mathcal{F}_W(\mathcal{O})$ , given  $\alpha_{\mathcal{U}_n}$ , the transformation  $\alpha_{\mathcal{U}_n} \alpha_{\mathcal{U}_n^a}^{-1}$ , with  $\mathcal{U}_n^a(\mathbf{x}) \equiv \mathcal{U}(\mathbf{x} - \mathbf{a})$ , is a Gauss gauge transformation and therefore it is implemented by a product  $V^{(a)}$  of Gauss operators  $V^{(a)}(\mathcal{U}^i)$ , which leave the vacuum vector invariant. Then, the following strong limit exists and defines an operator

$$S(\mathcal{U}_n) \equiv s - \lim_{|\mathbf{a}| \rightarrow \infty} V^{(a)},$$

which satisfies

$$S(\mathcal{U}_n) F S(\mathcal{U}_n)^{-1} = \alpha_{\mathcal{U}_n}(F), \quad \forall F \in \mathcal{F}_W, \quad S(\mathcal{U}_n) \Psi_0 = \Psi_0.$$

Clearly,  $C_{\mathcal{U}_n} \equiv V(\mathcal{U}_n) S(\mathcal{U}_n)^{-1}$  commutes with  $\mathcal{F}_W$ , belongs to the strong closure of  $\mathcal{F}_W$  and gives  $V(\mathcal{U}_n) F \Psi_0 = \alpha_{\mathcal{U}_n}(F) C_{\mathcal{U}_n} \Psi_0$ .

Since  $C_{\mathcal{U}_n}$  commutes with  $\mathcal{F}_W$ , they are completely characterized by their action on  $\Psi_0$ , where they only depend on the topological number  $n$ , commute with space time translations and satisfy eq. (5.2), because  $V(\mathcal{U}'_n)^{-1} V(\mathcal{U}_n)$ , and in particular  $V(\mathcal{U}_n^a)^{-1} V(\mathcal{U}_n)$ , are products of Gauss operators, which leave  $\Psi_0$  invariant.

Any open set contains an open cylinder  $\mathcal{O}$ , with base  $\mathcal{O}_0$ ,  $\mathcal{O} = \mathcal{O}_0 \times (t_1, t_2)$ , we denote by  $\mathcal{H}'(\mathcal{O})$  the subspace of vectors invariant under all the Gauss operators  $V(\mathcal{U}^\lambda)$ , with  $\text{supp } \mathcal{U}^\lambda \subset \mathcal{O}_0$  and by  $P(\mathcal{O})$

the corresponding projector, which belongs to (the strongly closed)  $\mathcal{F}_W(\mathcal{O})$ . Then, as in the proof of Proposition 3.2, one has that  $\forall \mathcal{U}$ ,  $V(\mathcal{U}^\lambda) V(\mathcal{U}) = V(\mathcal{U}) V((\mathcal{U}')^\lambda)$ , with  $\text{supp } (\mathcal{U}')^\lambda = \text{supp } \mathcal{U}^\lambda$ ; therefore

$$[V(\mathcal{U}), P(\mathcal{O})] = 0. \quad (5.4)$$

The operators  $C_n(\mathcal{O}) \equiv V(\mathcal{U}_n) P(\mathcal{O})$ , with  $\text{supp } \mathcal{U}_n \subset \mathcal{O}_0$ , only depend on the topological number  $n$ , since gauge transformation  $\mathcal{U}_n, \mathcal{U}'_n$  of the same homotopic class supported in  $\mathcal{O}_0$  differ by a product  $\mathcal{U}_G$  of Gauss transformations with the same support and  $V(\mathcal{U}_G) P(\mathcal{O}) = P(\mathcal{O})$ . This implies the first of eqs. (5.3).

Furthermore, since  $\forall \mathcal{U}$ ,  $V(\mathcal{U}) V(\mathcal{U}_n) = V(\mathcal{U}'_n) V(\mathcal{U})$ ,  $\text{supp } \mathcal{U}'_n \subset \mathcal{O}_0$ , by eq. (5.4), one has

$$V(\mathcal{U}) C_n(\mathcal{O}) = C_n(\mathcal{O}) V(\mathcal{U}),$$

i.e.  $C_n(\mathcal{O})$  are gauge invariant and therefore belong to  $\mathcal{A}(\mathcal{O})$ . Clearly, by construction  $C_n(\mathcal{O})$  commutes with  $\mathcal{A}$ .

Since  $P(\mathcal{O})\Psi = \Psi$ ,  $\forall \Psi \in \mathcal{H}'$ , and  $\cap_{\mathcal{O}} \mathcal{H}'(\mathcal{O}) = \mathcal{H}'$ , the remaining statements follow.

The unitary operators  $C_n$  are related to the topology of the gauge group, but they *do not* implement the (large) gauge transformations on the field algebra; they provide a unitary representation of the gauge group modulo the subgroup of Gauss transformations, through operators in the commutant, actually in the center, of  $\mathcal{F}_W$ .

The important feature of the operators  $C_n(\mathcal{O})$  is that of providing a representation of the group of gauge transformations localized in  $\mathcal{O}$ , modulo Gauss gauge transformations, through local operators which belong to the center  $\mathcal{Z}(\mathcal{O})$  of the observable algebra localized in  $\mathcal{O}$ . Thus, *the non-trivial topology of the gauge group is reflected by a non-trivial center of the local observable algebras  $\mathcal{A}(\mathcal{O})$ .*

The above structure implies that the non-trivial topology of the gauge group and eq. (4.1) force the breaking of chiral symmetry.

**Proposition 5.2** *Chiral symmetry  $\beta^\lambda$  is broken in any irreducible (or factorial) representation of the field algebra  $\mathcal{F}_W$ , defined by a Gauss invariant vacuum, as well as in any factorial (sub-)representation of the local observable algebra  $\mathcal{A}$  in the Gauss invariant subspace  $\mathcal{H}'$ .*

**Proof.** If the chiral symmetry  $\beta^\lambda$  is unbroken in  $\mathcal{H}$ , then there is a one-parameter group of unitary operators  $U^5(\lambda)$ ,  $\lambda \in \mathbf{R}$  satisfying

$$\beta^\lambda(F) = U^5(\lambda) F U^5(-\lambda), \quad \forall F \in \mathcal{F}_W, \quad U^5(\lambda) \Psi_0 = \Psi_0. \quad (5.5)$$

Such an action of  $U^5(\lambda)$  provides the unique strongly continuous extension of  $\beta^\lambda$  to  $\bar{\mathcal{F}}_W$ . Therefore, since the operators  $S(\mathcal{U}_n)$  are strong limits of Gauss operators, which are invariant under chiral transformations, by eqs. (4.1), (4.2), one has

$$\beta^\lambda(S(\mathcal{U}_n)) = S(\mathcal{U}_n).$$

On the other hand, by the localization of the  $V(\mathcal{U}_n)$ , eq. (4.2) applies and then eq. (4.1) (for  $R$  sufficiently large) gives

$$U^5(\lambda) V(\mathcal{U}_n) U^5(-\lambda) = \beta^\lambda(V(\mathcal{U}_n)) = e^{i2n\lambda} V(\mathcal{U}_n); \quad (5.6)$$

therefore

$$\beta^\lambda(C_n) = U^5(\lambda) C_n U^5(-\lambda) = e^{i2n\lambda} C_n. \quad (5.7)$$

This is inconsistent with an irreducible (or factorial) representation of  $\mathcal{F}_W$ , where the  $C_n$  are multiples of the identity.

Similarly, as a consequence of eqs. (4.1), (4.2), one has

$$\beta^\lambda(P_0) = U^5(\lambda) P_0 U^5(-\lambda) = P_0$$

and

$$\beta^\lambda(C_n P_0) = e^{i2n\lambda} C_n P_0. \quad (5.8)$$

Since  $C_n P_0$  is the limit of elements of the (strongly closed) local observable algebras  $\mathcal{A}(\mathcal{O})$ , it belongs to the center of the representation of  $\mathcal{A}$  in  $\mathcal{H}'$ ; this implies the instability of any factorial subrepresentation of  $\mathcal{A}$ , under  $\beta^\lambda$ .

The link between the non-trivial topology of the gauge group and the labeling of the factorial representations of the local observable algebra ( $\theta$  sectors) is clearly displayed in a (reducible) representation of the field algebra defined by a chirally invariant vacuum state. Such an invariance arises in an analysis based on the functional integral formulation and semiclassical considerations, [4] [5] as well as in rigorous treatments of soluble models (*in primis* the Schwinger model [19] [20] [21]).

In general, one obtains chirally invariant correlation functions by using chirally invariant boundary conditions in the functional integral in finite volume. [22] [1]

We shall therefore consider the case in which chiral symmetry is implemented in  $\mathcal{H}$  by a one-parameter group of unitary operators  $U^5(\lambda)$ , i.e. eqs. (5.5) hold.

**Proposition 5.3** *Under the above assumptions the factorial subrepresentations,  $\pi_\theta$ , of  $\mathcal{A}$  in  $\mathcal{H}'$  are labeled by an angle  $\theta$  ( $\theta$  sectors):  $\pi_\theta(C_n) = e^{i2n\theta}\mathbf{1}$ ,  $\theta \in [0, \pi)$ , (the corresponding groundstates are called  $\theta$  vacua).*

**Proof.** By eq. (5.2),  $C_n = C_1^n$  and by eq. (5.7) the spectrum of  $C_1$  is  $\sigma(C_1) = \{e^{i2\theta}; \theta \in [0, \pi)\}$ . This is also the spectrum of the operator  $C_1 P_0$  in  $\mathcal{H}'$ , by eq. (5.8).

The Hilbert space  $\mathcal{H}'$  has a central decomposition over the spectrum of  $C_1 P_0$  in  $\mathcal{H}'$ . Thus, one has

$$\mathcal{H}' = \int_{\theta \in [0, \pi)} d\theta \mathcal{H}_\theta, \quad C_n \mathcal{H}_\theta = e^{i2n\theta} \mathcal{H}_\theta,$$

since the spectral measure can be taken invariant under translations by eq. (5.7) and by the chiral invariance of the vacuum. By eq. (5.7),  $U^5(\alpha)$  intertwines between the sectors

$$U^5(\alpha) \mathcal{H}_\theta = \mathcal{H}_{\theta+\alpha} \quad (5.9)$$

and satisfies  $U^5(\pi) = (-1)^F$ , with  $F$  the fermion number ( $= 0$ , by our definition of  $\mathcal{F}_W$ ). Since the chiral symmetry commutes with time translations, the spectrum of the Hamiltonian is the same in all  $\theta$  sectors and “all the  $\theta$  vacua have the same energy”. The same decomposition applies to the representation of  $\mathcal{F}_W$  in  $\mathcal{H}$ .

Such a picture is exactly the same as in the quantum mechanical model of QCD structures discussed in Ref. [22]

Equation (5.6) provides a correct derivation of the equation

$$[V(\mathcal{U}_n), Q^5] = 2n, \quad (5.10)$$

which is at the basis of most of the standard discussions of chiral symmetry breaking in QCD. The standard derivation assumes that the

gauge transformations of the axial charge density extend to its space integral, giving the transformation properties of the chiral charge  $Q^5$ . Our derivation of the relation between chiral symmetry and large gauge transformations, eq. (5.6), does neither require the (usually assumed) convergence of the space integral of  $J_0^5$  to  $Q^5$ , nor that of its exponential, which are incompatible with Proposition 3.3.

It is worthwhile to stress that the breaking of chiral symmetry is governed by quite a different mechanism with respect to the Goldstone or the Higgs mechanism. In all the three cases the symmetry commutes with spacetime translations. However, in the Goldstone case, the symmetry breaking order parameter, typically an observable operator, has strong enough localization properties (preserved under time evolution) and its transformations under the symmetry are generated by a local conserved current. In the Higgs case, in positive gauges like e.g. the Coulomb gauge, the symmetry breaking order parameter is not an observable and it has a non-local time evolution, so that the (time independent) symmetry is not generated at all times by the associated conserved local Noether current.

In the axial  $U(1)$  case of QCD, contrary to statements appeared in the literature, the chiral transformations define a time independent symmetry of the observables. The Goldstone theorem, i.e. the presence of associated massless Goldstone bosons, is evaded by the impossibility of writing the corresponding symmetry breaking Ward identities, since the associated conserved Noether current does not exist, only its exponentials do (non-regular representation of the field algebra).

Actually, the chiral symmetry cannot be locally generated by unitary operators in any factorial representation of the observable algebra, because the local observable algebras have a center which is not left pointwise invariant under the chiral symmetry.

In conclusion, the non-regular Weyl quantization provides a strategy for putting the derivation of the vacuum structure and the chiral symmetry breaking in the temporal gauge of QCD in a more acceptable and convincing mathematical setting.



## 6 Regular temporal gauge

As discussed in the abelian case, one may look for an alternative realization of the temporal gauge, by weakening the condition of Gauss gauge invariance of the vacuum, so that the corresponding correlation functions of gauge dependent fields and not only those of their exponentials may be defined.

To be more precise, as before, one introduces a local field algebra  $\mathcal{F}$ , generated by  $\mathbf{A}(\mathbf{f})$ ,  $f_a^i \in \mathcal{S}(\mathbf{R}^4)$ , by the fermion fields, by their gauge invariant bilinears, by the axial current  $J_\mu^5$ , and by local operators  $V(\mathcal{U})$ , which implement the time independent gauge transformations  $\alpha_{\mathcal{U}}$ , eq. (2.7), represent the the group  $\mathcal{G}$  and satisfy

$$\alpha_{\mathcal{U}_n}(J_0(f_R \alpha_R)) = J_0(f_R \alpha_R) + 2n. \quad (6.1)$$

We denote by  $\mathcal{A}$  the gauge invariant (observable) subalgebra of  $\mathcal{F}$  and by  $V_G$  a generic monomial of the Gauss operators  $V(\mathcal{U}^\lambda)$ .

A **regular quantization of the temporal gauge** is defined by a (linear hermitian) vacuum functional  $\omega$  on  $\mathcal{F}$ , which is invariant under space-time translations and rotations and such that its restriction to the observable algebra  $\mathcal{A}$  satisfies positivity, Lorentz invariance and the relativistic spectral condition.

>From a constructive point of view, such a realization of the temporal gauge may be related to a functional integral quantization with a functional measure given by the Lagrangean of eq. (2.1) with the addition of the fermionic part (see Section 4). The invariance of the Lagrangean with respect to the residual gauge group after the gauge fixing  $\mathbf{A}^0 = 0$ , does not imply the corresponding residual gauge invariance of the correlation functions of  $\mathcal{F}$ , as discussed in the abelian case, [9]–[23] since an infrared regularization is needed which breaks the residual gauge invariance. Therefore, the Gauss constraint does not hold anymore.

The correlation functions of  $\mathcal{F}$  given by an  $\omega$  with the above properties define a vector space  $\mathcal{D} = \mathcal{F}\Psi_0$ , with  $\Psi_0$  the vector representing  $\omega$ , and an inner product on it  $\langle \cdot, \cdot \rangle$ , which is assumed to be left invariant by the operators  $V(\mathcal{U})$ .

It is further assumed that  $\omega$  satisfies the following **weak Gauss invariance**:

$$\omega(A V_G) = \omega(A), \quad \forall A \in \mathcal{A}, \quad \forall V_G,$$

equivalently

$$\langle A \Psi_0, V_G \Psi_0 \rangle = \langle A \Psi_0, \Psi_0 \rangle, \quad \forall A \in \mathcal{A}, \quad \forall V_G. \quad (6.2)$$

It follows that the vectors of the subspace  $\mathcal{D}'_0 \equiv \mathcal{A} \Psi_0$  are weakly Gauss invariant in the sense of eq.(6.2) and furthermore the space time translations  $U(a)$  leave  $\mathcal{D}'_0$  invariant. Thus,  $\omega$  defines a vacuum representations of  $\mathcal{A}$  in which the Gauss law holds.

The weak form of Gauss gauge invariance of the vacuum functional allows for the existence of the fields of  $\mathcal{F}$  as operators on  $\mathcal{D}$ , but the inner product cannot be semidefinite on  $\mathcal{D}$  (by the argument of Proposition 2.1). The subspace of vectors  $\Psi \in \mathcal{D}'_0$  with null inner product,  $\langle \Psi, \Psi \rangle = 0$ , is denoted by  $\mathcal{D}''_0$ .

Now, there is a substantial difference in the realization of the chiral symmetry, with respect to the representation defined by a Gauss invariant vacuum. Thanks to the weak form of the Gauss gauge invariance, the (smeared) conserved current  $J_\mu^5$  may be defined as an operator in  $\mathcal{D} = \mathcal{F} \Psi_0$  and the standard wisdom applies; in particular, for the infinitesimal variation  $\delta^5 F$  of the fields under chiral transformations, following Bardeen, one has

$$\delta^5 A = i \lim_{R \rightarrow \infty} [J_0^5(f_R \alpha_R), A], \quad \forall A \in \mathcal{A}. \quad (6.3)$$

In general, the representation  $\pi^{(0)}$  of the observable algebra defined by the vacuum vector  $\Psi_0$  may not be irreducible and therefore in order to discuss the breaking of the chiral symmetry one must decompose it into irreducible representations. Even if  $\omega(\delta^5 A) = 0$ , a symmetry breaking order parameter may appear in the irreducible components of  $\pi^{(0)}$ . Furthermore, such a decomposition of the vacuum functional on  $\mathcal{A}$  does not *a priori* extend to a decomposition of the vacuum expectations  $\langle \Psi_0, J_0^5(f_R \alpha_R) A \Psi_0 \rangle$ , since  $J_0^5(f_R \alpha_R)$  is not gauge invariant. Thus, one of the basic assumptions of the Goldstone theorem may fail and chiral symmetry breaking may not be accompanied by massless Goldstone bosons.

More definite statements can be made under the following reasonable assumption, hereafter referred to as the *existence of local implementers of the gauge transformations*:

- i) the subspace  $\mathcal{D}'$  generated by the vectors  $V(\mathcal{U})\mathcal{D}'_0$ , with  $\mathcal{U}$  running over  $\mathcal{G}$ , satisfies the weak Gauss constraint and semi-definiteness of the inner product,
- ii) if  $\text{supp } \mathcal{U} \subseteq \mathcal{O}$ , then,  $V(\mathcal{U})$  can be obtained as a “weak” limit of polynomials  $F_n$  of  $A_i^a$  and  $\psi$  localized in  $\mathcal{O}$ , in the following sense

$$\langle \Psi, V(\mathcal{U})\Phi \rangle = \lim_{n \rightarrow \infty} \langle \Psi, F_n \Phi \rangle, \quad \forall \Psi, \Phi \in \mathcal{D}. \quad (6.4)$$

Property i) is supported by the fact that the states defined by the vectors  $V(\mathcal{U})A\Psi_0$ ,  $A \in \mathcal{A}$ , are weakly Gauss invariant and positive; in fact  $\forall A, B, C \in \mathcal{A}$ ,

$$\begin{aligned} \langle AV(\mathcal{U})B\Psi_0, V_G V(\mathcal{U})C\Psi_0 \rangle &= \langle \alpha_{\mathcal{U}}(A)B\Psi_0, V'_G C\Psi_0 \rangle = \\ &= \langle AB\Psi_0, C\Psi_0 \rangle = \langle AV(\mathcal{U})\Psi_0, V(\mathcal{U})C\Psi_0 \rangle. \end{aligned}$$

The stability under  $V(\mathcal{U})$  of a weakly Gauss invariant subspace, which includes  $\mathcal{D}'_0$ , is automatically satisfied if such a subspace may be selected by a gauge covariant subsidiary condition. Weak Gauss invariance of  $\mathcal{D}'$  is also implied by the following stronger form of the weak Gauss invariance of the vacuum functional

$$\omega(AV(\mathcal{U}^\lambda)V(\mathcal{U})) = \omega(AV(\mathcal{U})), \quad \forall A \in \mathcal{A}, \quad \forall \mathcal{U}^\lambda, \mathcal{U}. \quad (6.5)$$

Property ii) is supported by the localization of the gauge transformations so that the  $V(\mathcal{U})$  are local relative to the field algebra, with localization region given by the support of the corresponding gauge transformation.

The fields  $F$  which leave  $\mathcal{D}'$  invariant also leave the subspace  $\mathcal{D}''$  of null vectors of  $\mathcal{D}'$  invariant and therefore define unique gauge invariant operators  $\hat{F}$  in the “physical” quotient space  $\mathcal{D}_{phys} \equiv \mathcal{D}'/\mathcal{D}''$ , which is the analog of the Gauss invariant subspace  $\mathcal{H}'$  of the non-regular realization of the temporal gauge. Thus, to all effects such fields can be considered as observable fields; in the following we shall take as *observable algebra localized in  $\mathcal{O}$* ,  $\hat{\mathcal{A}}(\mathcal{O})$ , the algebra of operators in  $\mathcal{D}_{phys}$  generated by fields localized in  $\mathcal{O}$  which leave  $\mathcal{D}'$  invariant and as observable algebra  $\hat{\mathcal{A}} \equiv \cup_{\mathcal{O}} \hat{\mathcal{A}}(\mathcal{O})$ . [24] In particular, the local operators  $V(\mathcal{U}_n(\mathcal{O}))$  are weakly gauge invariant and therefore they define unique operators  $\hat{T}_{\mathcal{U}_n(\mathcal{O})} \in \hat{\mathcal{A}}(\mathcal{O})$  in  $\mathcal{D}_{phys}$ .

By the same arguments discussed before, the  $\hat{T}_{\mathcal{U}_n}$  depend only on  $n$ , are invariant under space time translations and satisfy

$$\hat{T}_n \hat{T}_m = \hat{T}_{n+m}, \quad \hat{T}_0 = \mathbf{1}. \quad (6.6)$$

Moreover, since one may write  $V(\mathcal{U}_n) = V(\mathcal{U}_n^a) V_G$ , for any local  $F$ , which leaves  $\mathcal{H}'$  invariant, one has

$$\hat{T}_n \hat{F} = \hat{F} \hat{T}_n. \quad (6.7)$$

This implies that the  $\hat{T}_n$  generated an abelian group  $\mathcal{G}_T$  and belong to the center  $\mathcal{Z}(\mathcal{O})$  of  $\hat{\mathcal{A}}(\mathcal{O})$ ,  $\forall \mathcal{O}$ .

Furthermore, the local generation of the infinitesimal chiral transformations, eq. (6.3), implies weak continuity of the derivation  $\delta^5$  on the local field algebras  $\mathcal{F}(\mathcal{O})$  and by property ii), the infinitesimal chiral transformations of the local implementers  $V(\mathcal{U}_n)$  of the large gauge transformations are determined by eq. (6.1), i.e.

$$\begin{aligned} \langle \mathcal{D}, \delta^5(V(\mathcal{U}_n) \mathcal{D}) \rangle &= \lim_{m \rightarrow \infty} \langle \mathcal{D}, \delta^5(F_m) \mathcal{D} \rangle = \\ \lim_{m \rightarrow \infty} i \langle \mathcal{D}, [J_0^5(f_R \alpha_R), F_m] \mathcal{D} \rangle &= i \langle \mathcal{D}, [J_0^5(f_R \alpha_R), V(\mathcal{U}_n)] \mathcal{D} \rangle = \\ &= i 2n \langle \mathcal{D}, V(\mathcal{U}_n) \mathcal{D} \rangle, \end{aligned} \quad (6.8)$$

for  $R$  sufficiently large so that  $f_R(\mathbf{x}) = 1$  on the localization region of  $\mathcal{U}_n$ . Thus, one has

$$\delta^5(\hat{T}_n) = i 2n \hat{T}_n, \quad (6.9)$$

and if the chiral symmetry is unitarily implemented in  $\mathcal{D}_{phys}$  the observable algebra (in  $\mathcal{D}_{phys}$ ) has a non-trivial center  $\mathcal{Z}$ .

**Proposition 6.1** *Under the above general assumptions, one has*

- i) the non-trivial topology of the gauge group gives rise to a center of the observable algebra (in the physical space  $\mathcal{D}_{phys}$ ), which not left pointwise invariant under the chiral symmetry,*
- ii) the chiral symmetry is broken in any factorial representation of the observable algebra,*
- iii) the decomposition of the physical Hilbert space  $\mathcal{H}_{phys} \equiv \overline{\mathcal{D}_{phys}}$  over the spectrum of  $\hat{T}_1$  defines representations of the observable algebra labeled by an angle  $\theta \in [0, \pi)$ , giving rise to the  $\theta$  vacua structure,*

iv) the expectations  $\omega_\theta(J_0^5(f_R\alpha_R)A)$ ,  $A \in \mathcal{A}$ , with  $\omega_\theta$  invariant under gauge transformations, cannot be defined and a crucial condition of the Goldstone theorem fails.

**Proof.** Most of the arguments are essentially the same as in the non-regular realization. In particular, an unbroken chiral symmetry in a factorial representation of the algebra of observable is incompatible with the non-trivial chiral transformations of its center.

By eq.(6.9), the spectrum of  $\hat{T}_1$  is  $\{e^{i2\theta}, \theta \in [0, \pi)\}$ , and, even if  $J_0^5$  is well defined as an operator in  $\mathcal{D}$ , the existence of the expectations  $\omega_\theta(J_0^5(f_R\alpha_R)A)$  would lead to the same inconsistency as in eq. (4.4).

## 7 The Schwinger model in the temporal gauge

The general features discussed above are exactly reproduced by the Schwinger model in the temporal gauge, usually regarded as a prototype of the non-perturbative QCD structures; in particular, the assumptions about the local implementers of the large gauge transformations hold.

The bosonized Schwinger model in the temporal gauge is formally described by the following Lagrangean density

$$\mathcal{L} = \frac{1}{2}(\partial_0\varphi)^2 - \frac{1}{2}(\partial_1\varphi)^2 + \partial_0\varphi A_1 + \frac{1}{2}(\partial_0 A_1)^2, \quad (7.1)$$

where  $\varphi$  is the pseudoscalar field which bosonizes the fermion bilinears and therefore is an angular variable, and  $A_1$  is the gauge vector potential.

The time evolution is formally determined by the following canonical equations

$$\pi = \partial_0\varphi + A_1, \quad \partial_0 A_1 = E, \quad \partial_0\varphi = \Delta\varphi, \quad \partial_0 E = \partial_0\varphi. \quad (7.2)$$

### 1) Representation by a Gauss invariant vacuum

As exponential field algebra, we take the algebra generated by the unitary operators

$$V_\varphi(f), \quad \int dx f = n, \quad V_A(h), \quad V_E(g), \quad V_\pi(g), \quad f, g, h, \in \mathcal{D}(\mathbf{R}), \quad (7.3)$$

formally corresponding to the exponentials  $e^{i\varphi(f)}$ ,  $e^{iA_1(h)}$ ,  $e^{iE(g)}$ ,  $e^{i\pi(g)}$ , respectively, and satisfying the Weyl commutation relations, with above restriction on  $f$ , required by the periodicity of  $\varphi$ .

The time independent gauge transformations

$$\alpha_{\mathcal{U}}(V_A(h)) = V_A(h + \mathcal{U}\partial_1\mathcal{U}^{-1}), \quad \mathcal{U}(x_1) - \mathbf{1} \in \mathcal{D}(\mathbf{R}),$$

$$\alpha_{\mathcal{U}}(V_{\pi}(g)) = V_{\pi}(g + \mathcal{U}\partial_1\mathcal{U}), \quad (7.4)$$

$\varphi$  and  $E$  being left invariant, are generated by the local operators

$$V(\mathcal{U}) \equiv V_{\varphi}(f) V_E(-f) \equiv V(f), \quad f = \mathcal{U}\partial_1\mathcal{U}^{-1}, \quad \int dx f = n. \quad (7.5)$$

The gauge functions  $f$  with  $\int dx f = 0$ , i.e. those of the form  $f = \partial_1 g$ ,  $g \in \mathcal{D}(\mathbf{R})$ , define the Gauss transformations and those with  $\int dx f = n$ ,  $n \neq 0$ , define the large gauge transformations and shall be labeled by the topological number  $n$ . Clearly, if  $\mathcal{U}$  is localized in  $\mathcal{O}$ , equivalently  $\text{supp } f \subseteq \mathcal{O}$ , then  $V(f) \in \mathcal{F}_W(\mathcal{O})$ , so that our assumption of local implementability is verified. The dynamics is defined by eqs. (7.2) and therefore  $e^{i\sigma(f)} \equiv e^{i(\varphi-E)(f)}$  is independent of time.

The chiral transformations  $\beta^{\lambda}$  are defined by

$$\beta^{\lambda}(V_{\varphi}(f_n)) = e^{i2n\lambda} V_{\varphi}(f_n), \quad (7.6)$$

all the other exponential fields being left invariant. Thus, as argued in general, the anomaly of the gauge invariant axial current  $j_{\mu}^5 = \partial_{\mu}\varphi$ ,  $\partial^{\mu}j_{\mu}^5 = \varepsilon_{\mu\nu}\partial^{\mu}A^{\nu}$  does not prevent the chiral symmetry from defining a one-parameter group of automorphisms of the (exponential) field algebra and of its gauge invariant subalgebra  $\mathcal{A}$ , locally generated by the unitary operators  $V_R^5(\lambda) \equiv V_{\pi}(f_R\alpha_R)$ .

The GNS representation of  $\mathcal{F}_W$  by a Gauss invariant state  $\omega$  is characterized by a representative vector  $\Psi_0$  which satisfies

$$V(\partial_1 g) \Psi_0 = \Psi_0, \quad g \in \mathcal{D}(\mathbf{R}). \quad (7.7)$$

The Gauss invariance of the vacuum vector is independently required by the condition of positivity of the energy, using the positivity of the state  $\omega$  and the invariance under space translations. In fact, since  $V(f)$

commutes with the Hamiltonian, one can take  $\Psi_0$  as an eigenstate of  $V(f)$ , i.e.  $V(f)\Psi_0 = e^{i\lambda(f)}\Psi_0$ ; then, by introducing

$$V(t) \equiv e^{i\alpha_t(\pi(g) - \Delta A(g))} = V(0)e^{it\Delta\sigma(g) + it \int dx \Delta(1-\Delta)g^2},$$

one gets

$$\begin{aligned} (V(0)\Psi_0, H V(0)\Psi_0 &= i(d/dt)(V(0)\Psi_0, V(t)\Psi_0)_{t=0} = \\ &= -\lambda(\Delta g) - \int dx \Delta(1-\Delta)g^2. \end{aligned}$$

Therefore, the positivity of the energy  $\forall g$  requires that the functional  $\lambda(\Delta g)$  be  $\leq 0$ ,  $\forall g$ , and therefore  $= 0$ , since it is linear in  $g$ . On the other hand, since any  $f$  can be decomposed as the sum  $h_1 \int dx f + h_2 \int dx x f + \Delta h_3$ , with  $h_i \in \mathcal{D}(\mathbf{R})$ , the invariance under space translations requires  $\lambda(h_2) = 0$  and one gets  $\lambda(f) = 0$ ,  $\forall f = \partial_1 g$ .

By the same argument of Proposition 2.1, the Gauss invariance of the vacuum vector implies the vanishing of all the expectations  $\omega(FV_A(h))$ , with  $F$  any element of the gauge invariant subalgebra  $\mathcal{A}$  of  $\mathcal{F}_W$ , unless  $h = 0$ . Hence, we are left with the correlation functions of the gauge invariant fields.

Since eqs. (7.2) imply  $\square E + E = \Delta\sigma$ , and  $E$  is a pseudoscalar field, the vacuum correlations functions of  $E$  are those of a free pseudoscalar field of mass  $= 1$ . By the Gauss invariance of the vacuum, the expectations  $\omega(V(f_n) \alpha_t(V_E(g)))$  depend on  $f_n$  only through the topological number  $n$  and therefore define operators  $T_n$ , which are invariant under spacetime translations and satisfy  $T_n T_m = T_{n+m}$ .

The residual arbitrariness is therefore that of the representation of the abelian algebra  $G_T$ , generated by the operators  $T_n$  in the subspace  $\mathcal{A}\Psi_0$ . The  $\theta$  vacua are characterized by the expectations

$$\omega_\theta(V(f_n) A) = \omega_\theta(T_n A) = e^{i2n\theta} \omega_\theta(A), \quad \forall A \in \mathcal{A}. \quad (7.8)$$

On the other hand, the reducible representation defined by a chirally invariant vacuum is characterized by the expectations

$$\omega(V(f_n) A) = \omega(T_n A) = \delta_{n0} \omega(A), \quad \forall A \in \mathcal{A}, \quad \beta^\lambda(A) = A.$$

It is easy to check that all the general features of the QCD case discussed in Sections 3-5, in particular the evasion of the Goldstone

theorem, the breaking of chiral symmetry in any irreducible or factorial representation of the observable algebra, as a consequence of the non-trivial topology of the gauge group and the  $\theta$  vacua structure are exactly reproduced.

## 2) Regular representation

As local field algebra  $\mathcal{F}$  we take the canonical algebra generated by the fields  $A_1(h)$ ,  $E(g)$ ,  $V_\varphi(f)$ , with  $\int dx f = n$ ,  $\partial_0 \varphi(f)$  with the equal time commutation relations

$$[A_1(x_1, t), E(y_1, t)] = i\delta(x_1 - y_1),$$

$$[V_\varphi(f), (\partial_0 \varphi + A_1)(g)] = i \int dx_1 f g V_\varphi(f).$$

The euclidean functional integral corresponding to the Lagrangean of eq. (7.1) yields well defined correlation functions of  $E$  and  $\partial_\mu \varphi$  satisfying the (weak) Gauss law constraint. As before, the two-point function  $W_E(x)$  of  $E$  is that of free pseudoscalar massive field. The correlation functions of  $e^{i\varphi}$  involve a zero mode  $\varphi_0$  and crucially depend on the boundary conditions (in finite volume). Any boundary condition satisfying positivity yields the (weak) Gauss law holds for the expectations of all the gauge invariant variables. Periodic boundary conditions in finite volume give chiral invariant correlation functions and therefore for any polynomial function  $\mathcal{P}$ ,

$$< e^{i\varphi_0} \mathcal{P}(\partial_1 \varphi, E) > = \delta_{n0} < \mathcal{P}(\partial_1 \varphi, E) >,$$

(for a general discussion of the role of the boundary conditions in QCD and in related models see Ref. [25] [1] [22]).

An infrared subtraction is needed for defining the correlation functions of  $A_1$ . The most general two-point function  $W_A$  of  $A_1$  must satisfy  $-(d^2/dt^2)W_A(x) = W_E(x)$  and therefore has the following form ( $\omega(k_1) \equiv \sqrt{k_1^2 + 1}$ )

$$\frac{1}{2\pi} \int dk_1 \omega(k_1)^{-3} e^{i(\omega(k_1)x_0 - k_1 x_1)} - \frac{1}{2}i (\delta(x_1) - \frac{1}{2}e^{-|x_1|})x_0$$

$$+ B(x_1)x_0 + C(x_1). \quad (7.9)$$



Locality requires  $B(x_1) = -B(-x_1)$ ,  $C(x_1) = C(-x_1)$ . The term linear in  $x_0$  violate positivity. As discussed in the QED case [9] the function  $C$  can be removed by an operator time independent gauge transformation and  $B = 0$  if  $w$  is invariant under the  $CP$  symmetry  $\Gamma$ :  $\Gamma(\varphi(x_1)) = \varphi(-x_1)$ ,  $\Gamma(A_1(x_1)) = A_1(-x_1)$ .

The two-point function  $\langle \partial_1 \varphi(x_1, 0) A_1(y_1, 0) \rangle$  parametrizes the infrared regularization of the functional integral corresponding to the Lagrangean of eq.(7.1) and can be taken to vanish. Then, all the two-point functions involving  $\partial_1 \varphi$  and  $A_1$  are determined. The corresponding  $n$ -point-functions can be taken as factorized (Gaussian).

Then, we are left with the correlation functions involving the zero mode  $\varphi_0$ , equivalently the correlation functions involving  $e^{i\sigma(f_n)}$ ,  $n \neq 0$ .

By the weak Gauss invariance, the one-point function  $\langle e^{i\sigma(f_n)} \rangle_0 \equiv s_n$  only depends on the topological number and

$$s_n = \bar{s}_{-n}, \quad \langle e^{i\sigma(f_n)} e^{i\sigma(f_m)} \rangle = \langle e^{i\sigma(f_n+f_m)} \rangle = s_{n+m}.$$

Semi-definiteness of the subspace  $\mathcal{A}\Psi_0 = \mathcal{D}'$  implies that the sequence  $\{s_n\}$  is of positive type and (since  $e^{i\sigma(f_n)}$  commutes with  $\mathcal{A}$ ) the vacuum functional on  $\mathcal{A}$  has the decomposition

$$w(A) = \int_0^\pi d\mu(\theta) \omega_\theta(A), \quad \forall A \in \mathcal{A}, \quad (7.10)$$

$$\omega_\theta(e^{i\sigma(f_n)} A) = e^{i2n\theta} w(A), \quad \forall A = \mathcal{P}(\partial_1 \varphi, E).$$

The representation of  $\mathcal{A}$  is the same as in positive case; in fact, it only depends on the weak Gauss invariance of the vacuum.

The (non-positive) extension to the gauge field algebra  $\mathcal{F}$  is given by the correlation functions  $\langle e^{i\sigma(f_n)} A_1(z_1) \dots A_1(z_k) \rangle$ ,  $z_i = (x_1^{(i)}, x_0^{(i)})$ . For simplicity, we consider the case of a chirally invariant vacuum functional  $w$ . Then, all such correlation functions for  $n \neq 0$  vanish and  $s_n = \delta_{n0}$ , corresponding to  $d\mu(\theta) = d\theta/\pi$ .

In agreement with the general analysis of Sections 3,4, the chiral symmetry cannot be locally generated in the physical space  $\mathcal{D}_{phys} = \mathcal{D}'/\mathcal{D}''$ ; in particular, the density of the axial current  $J_0^5 = \partial_0 \varphi + A_1$ , which generates the symmetry on  $\mathcal{F}$ , cannot be defined there, by the argument of Proposition 4.1. Thus, the breaking of the chiral symmetry in any factorial representation of the observable algebra does not require the existence of massless Goldstone bosons.

Since the correlation functions  $\langle e^{i\sigma(h)} \mathcal{P}(\partial_1 \varphi, E, A_1) \rangle$  factorize in terms of two-point functions, which satisfy the cluster property, the limits of the correlation functions of  $e^{i\sigma(f_n^a)}$ ,  $f_n^a(x_1) \equiv f_n(x_1 - a)$ , when  $|a| \rightarrow \infty$ , exist and define the analogs of the operators  $C_n$  of Section 5. Then, by the chiral invariance of  $w$ , one has  $w(C_n) = \delta_{n0}$  and  $w$  may be decomposed as a direct integral of (indefinite) functionals  $w_\theta$  on  $\mathcal{F}$ , characterized by the expectations  $w_\theta(C_n) = e^{i2n\theta}$ , which do not lead to a decomposition in which the  $C_n$  are multiples of the identity. Clearly,  $w_\theta$  coincides with the  $\theta$  vacuum on  $\mathcal{A}$ :  $w_\theta(A) = \omega_\theta(A)$ ,  $\forall A \in \mathcal{A}$ , but represents a non-positive extension to  $\mathcal{F}$ , which is not invariant under the gauge transformations.

The corresponding chiral symmetry breaking Ward identities

$$w_\theta([J_R^5, e^{i\sigma(f_n)}]) = i2n w_\theta(e^{i\sigma(f_n)}) = i2n e^{i2n\theta}, \quad J_R^5 \equiv J_0^5(f_R \alpha_R),$$

involve correlation functions  $\langle J_0^5(x) e^{i\sigma(f_n)} \rangle$  which are independent of time, but the time independent vectors, playing the role of the Goldstone bosons, are equivalent to  $e^{i2n\theta} \Psi_\theta$  in the physical space  $\mathcal{H}_\theta$  and therefore do not give rise to zero energy states different from the vacuum. Actually, the possibility of writing a chiral symmetry breaking Ward identity in terms of a commutator with a current operator in the physical space (with the vacuum vector  $\Psi_\theta$  in their domain) is excluded, since any such a commutator with  $e^{i\sigma(f_n)}$  has vanishing vacuum expectation.

It is easy to check that all the general features and assumptions of the QCD case are realized.

## References

- [1] B. Bock-Bavnbek, G. Morchio, F. Strocchi and K.P. Wojciechowski, Jour. Geom. Phys., **22**, 219 (1997)
- [2] G. t' Hooft, Phys. Rev. Lett. **37**, 8 (1976)
- [3] R. Jackiw and C. Rebbi, Phys. Rev. Lett. **37**, 172 (1976)
- [4] C.G. Callan, R.F. Dashen and D. Gross, Phys. Lett. **B 63**, 34 (1976)
- [5] S. Coleman, *Aspects of Symmetry*, Cambridge Univ. Press 1985, Chapter 7, Sect.3
- [6] S. Weinberg, *The Quantum Theory of Fields. Vol. II. Modern Applications*, Cambridge University Press 1996, Sect. 23.6
- [7] See the excellent lectures by R. Jackiw, Topological investigations of quantized gauge theories, in S.B. Treiman, R. Jackiw, B. Zumino and E. Witten, *Current Algebra and Anomalies*, World Scientific 1985, pp. 211-359
- [8] W.A. Bardeen, Nucl. Phys. **B 75**, 246 (1974)
- [9] J. Löffelholz, G. Morchio and F. Strocchi, Jour. Math. Phys. **44**, 5095 (2003)
- [10] A. Bassetto, G. Nardelli and R. Soldati, *Yang-Mills Theories in Algebraic Non-covariant Gauges*, World Scientific 1991
- [11] F. Acerbi, G. Morchio and F. Strocchi, Lett. Math. Phys. **27**, 1 (1993); J. Math. Phys. **34**, 899 (1993)
- [12] B. Schroer and P. Stichel, Comm. Math. Phys. **3**, 258 (1966)
- [13] H.D. Maison, Nuovo Cim. **11A**, 389 (1972)
- [14] M. Requardt, Comm. Math. Phys. **50**, 259 (1976)
- [15] G. Morchio and F. Strocchi, Jour. Math. Phys. **44**, 5569 (2003)

- [16] G. Morchio and F. Strocchi, Jour. Phys. A: Math. Theor. **40**, 3173 (2007)
- [17] R.J. Crewther, Chiral Properties on Quantum Chromodynamics, in *Field Theoretical Methods in Particle Physics*, W. Rühl ed., Reidel 1980, pp. 529-590
- [18] The existence of local implementers of a gauge symmetry has been proved in general for local covariant field algebras, by exploiting the Reeh-Schlieder property; see D. Buchholz, S. Doplicher, R. Longo and J.E. Roberts, Rev. Math. Phys. **4**, 49 (1992)
- [19] A.Z. Capri and R. Ferrari, Nuovo Cim. **A62**, 273 (1981); Jour. Math. Phys. **25**, 141 (1983)
- [20] G. Morchio, D. Pierotti and F. Strocchi, Ann. Phys. **188**, 217 (1988)
- [21] F. Strocchi, *Selected Topics on the General Properties of Quantum Field Theory*, World Scientific 1993, Sect. 7.4
- [22] J. Löffelholz, G. Morchio and F. Strocchi, Ann. Phys. **250**, 367 (1996)
- [23] J. Löffelholz, G. Morchio and F. Strocchi, Groud state and functional integral representation of the CCR algebra with free evolution, arXiv math-ph/0212037.
- [24] For the discussion of such weak gauge invariance and its characterization of the observables in indefinite quantum field theories, see F. Strocchi and A.S. Wightman, Jour. Math. Phys. **15**, 2198 (1974)
- [25] G. Morchio and F. Strocchi, Boundary Terms, Long Range Effects, and Chiral Symmetry Breaking, Lectures at the XXIX Int. Universitätswochen Schladming, March 1990, in *Fields and Particles*, H. Mitter and W. Schweiger eds., Springer 1990, pp.171-214.